

AD-A111 215

ROCHESTER UNIV NY DEPT OF CHEMISTRY

F/6 20/10

QUANTUM MODEL OF SELECTIVE MULTIPHOTON PROCESSES: THE ROLE OF E--ETC(U)

JAN 82 J LIN, T F GEORGE

N00014-80-C-0472

UNCLASSIFIED

TR-17

NL

OF /
AD
A11215




END
DATE
FILMED
82
DTIC

CLASSIFIED

1 OF 1

AD

A/11215



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A111215

OFFICE OF NAVAL RESEARCH

Contract N00014-80-C-0472

Task No. NR 056-749

TECHNICAL REPORT No. 17

Quantum Model of Selective Multiphoton
Processes: The Role of External (Field) and
Internal (Anharmonic Coupling) Detunings

by

Jui-teng Lin and Thomas F. George

Prepared for Publication

in

Theoretica Chimica Acta

University of Rochester
Department of Chemistry
Rochester, New York 14627

January, 1982

Reproduction in whole or in part is permitted for any purpose
of the United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
SELECTED
FEB 22 1982
A

DTIC FILE COPY

82 02 22 112

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 17	2. GOVT ACCESSION NO. 40 A111 215	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Quantum Model of Selective Multiphoton Processes: The Role of External (Field) and Internal (Anharmonic Coupling) Detunings		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
7. AUTHOR(s) Jui-teng Lin and <u>Thomas F. George</u>		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Rochester Department of Chemistry Rochester, New York 14627		8. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0472
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program Code 472 Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 056-749
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE January 1982
		13. NUMBER OF PAGES 16
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for publication in Theoretica Chimica Acta, in press.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) MULTIPHOTON EXCITATION LASER DETUNING INFRARED LASER ANHARMONIC COUPLING ENERGY TRANSFER HOMOGENEOUS SYSTEM (SF ₆) INTRAMOLECULAR RELAXATION HETEROGENEOUS SYSTEM (H/W)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Quantum equations of motion describing the energy transfer dynamics via intramolecular anharmonic coupling are presented and solved numerically. Dynamical features of the average excitations of a homogeneous system (SF ₆) and a heterogeneous system (H/W) are quantitatively discussed in terms of the intramolecular vibrational relaxation (IVR) rates. Possible mechanisms of surface-enhanced desorption via low-power laser radiation are proposed.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601


Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Quantum Model of Selective Multiphoton Processes:
The Role of External (Field) and
Internal (Anharmonic Coupling) Detunings*

Jui-teng Lin and Thomas F. George[†]
Department of Chemistry
University of Rochester
Rochester, New York 14627 USA

Quantum equations of motion describing the energy transfer dynamics via intramolecular anharmonic coupling are presented and solved numerically. Dynamical features of the average excitations of a homogeneous system (SF_6) and a heterogeneous system (H/W) are quantitatively discussed in terms of the intramolecular vibrational relaxation (IVR) rates. Possible mechanisms of surface-enhanced desorption via low-power laser radiation are proposed.



1. Introduction

The selective nature of laser-induced processes, which crucially depend on the intramolecular vibrational relaxation (IVR) rate and on the pumping rate of the laser radiation, have been widely studied for homogeneous (gas-phase) systems¹⁻³ and more recently for heterogeneous systems.⁴⁻⁷

For homogeneous systems, several types of selectivity have been discussed in terms of the relative magnitudes of the pumping rate, the IVR rate, the intermolecular vibration-vibration and vibration-translation rates: (1) mode selective (for fast pumping), (2) molecule-selective (for fast IVR rate), (3) nonselective vibrational excitation (for strong intermolecular vibration-vibration coupling) and (4) thermal excitation of the system.¹

For heterogeneous systems, e.g., molecules (atoms) adsorbed on a solid surface, the nature of the laser selective effects are characterized not only by the intra- and intermolecular relaxation processes of the adsorbed species and the pumping rate of the laser field, but also by the surface-induced damping factor (via multiphonon processes) and the laser-induced surface dynamical phenomena, e.g., migration, desorption and predissociation.⁸⁻¹¹ The laser-stimulated surface processes then may be: (1) mode selective, (2) adspecies selective, (3) migration (lattice-site) selective, (4) local heating and (5) nonselective thermal heating.⁸

In a homogeneous system, the mode- and/or molecule-selective excitations are possible by a high-power short pulse when the IVR rate is small.^{1,12} For a heterogeneous system, the laser-

stimulated selective surface bond breaking (desorption) is also possible when the phonon-induced damping rate is small for high-order multiphonon processes.^{5,13} The IVR rate therefore plays the essential role in the nature of the laser-selective excitations, both for homogeneous and heterogeneous systems. In this Letter, the dynamical features of the intramolecular modes governed by the intramolecular anharmonic couplings are studied for two systems: (A) SF₆ (gas phase) and (B) H/W (hydrogen atom adsorbed on a tungsten surface).

2. Quantum Equations of Motion for a Nonlinear Open System

The total Hamiltonian describing an anharmonic quantum oscillator subject to laser (IR) radiation of an open system may be written as

$$H(t, Q_1, Q_2, \dots, Q_N) = H_0^{\text{eff}} + \lambda H_{\text{anh}} + H'(t), \quad (1)$$

where H_0^{eff} is the effective unperturbed Hamiltonian of the system (with N normal modes), λH_{anh} is the intramolecular anharmonic coupling and $H'(t)$ is the laser field interaction Hamiltonian. In a second-quantization representation, the individual terms of the total Hamiltonian may be expressed by

$$H_0^{\text{eff}} = \sum_j \hbar \omega_j a_j^\dagger a_j - i \sum_j \hbar \Gamma_j / 2, \quad (2.a)$$

$$H_{\text{anh}} = \sum_{i+j+k} \sum_{l,m,n} \left[F_{lmn}^{ijk} (a_i^\dagger)^l (a_j^\dagger)^m (a_k^\dagger)^n + G_{lmn}^{ijk} (a_i^\dagger)^l (a_j^\dagger)^m (a_k)^n \right] + \dots, \quad (2.b)$$

$$H'(t) = \sum_j V_j(t) (a_j^\dagger + a_j). \quad (2.c)$$

In eq. (2.a), the imaginary part of the complex effective Hamiltonian Γ_j takes account of the level broadening of the open system due to factors such as collisional (pressure) broadening for a homogeneous system or surface-phonon-induced damping for a heterogeneous system. In eq. (2.b), two typical terms of the anharmonic coupling are shown: the first term representing T_2 phase relaxation processes, where no vibrational energy among the intramolecular modes is exchanged, and the second term representing the T_1 energy relaxation processes where the photon energy is deposited in the active mode(s) and populated in all the other inactive mode via anharmonic coupling. In eq. (2.c), $V_j(t) \propto V_{j0} \cdot \cos \omega t$ is proportional to the derivative of the dipole moment and the effective local electric field of the j -th mode with laser frequency ω . We note that the anharmonic coupling factors F_{lmn}^{ijk} and G_{lmn}^{ijk} are related to the p -th derivatives of the anharmonic potential energy (with respect to the normal coordinates) with $p = l+m+n$ being the order of the multiquantum coupling.

In order to study the energy transfer among the intramolecular modes, we shall first set up the equations of motion of the energy-related relevant quantities. The solution of the time-dependent Schrödinger equation

$$i\hbar[\partial\Psi/\partial t] = H\Psi, \quad (3)$$

with the total Hamiltonian H given by eq. (1) may be written as¹⁴

$$\Psi(t) = \exp\left\{-\frac{i}{\hbar} H_0^{\text{eff}} t + A(t) + \sum_j \left(B_j(t) a_j^\dagger - B_j^*(t) a_j \right)\right\} \Psi(0). \quad (4)$$

where $A(t)$ and $B(t)$ are time-dependent coefficients.

Substituting eq. (4) into eq. (3) with the expression of the total Hamiltonian [eqs. (1) and (2)], we find the equations of motion of the amplitude functions B_j ($j=1,2,\dots,N$) to be

$$i\hbar \frac{dB_j}{dt} = \lambda \sum_{i+j+k} \sum_{lmn} m \left\{ F_{lmn}^{ijk} (B_i^* B_j)^l (B_j^*)^{m-1} (B_k B_k)^n + \right. \\ \left. G_{lmn}^{ijk} (B_i^*)^l (B_j^*)^{m-1} (B_k)^n \exp(i\Delta_I t) \right\} \\ + (V_{j0}/2) \exp(i\Delta_E^j t) - (i\Gamma_j/2) B_j, \quad (5)$$

where the external detuning (Δ_E^j) and the internal detuning (Δ_I) are defined by $\Delta_E^j = 2\pi(\nu_{j+1} - \nu_j) - \omega$ and $\Delta_I = 2\pi(\ell\nu_i + m\nu_j - n\nu_k)$. In deriving eq. (5), we have used the usual rotating-wave approximation and operator algebra such as¹⁴

$$[\exp(B^* a)] F(a^\dagger) [\exp(-B^* a)] = F(a^\dagger + B_j^*), \quad (6)$$

where $F(a^\dagger)$ is any operator function and B_j are c-numbers.

The important features of the quantum equations of motion [eq. (5)] are: (1) the j -th normal mode of the system is strongly coupled to the laser field when $V_{j0} \neq 0$ and the external detuning is not far-off the resonance condition $\Delta_E^j = 0$ (due to the anharmonicity, we shall find later that the optimal detuning is red-shifted, i.e., $\Delta_E^j > 0$); (2) the intramolecular coupling is governed by the coupling strengths λF_{lmn}^{ijk} , λG_{lmn}^{ijk} and the internal detuning Δ_I : for the intramolecular coupling to be significant, we shall expect a near

resonance $\Delta_I \approx 0$, which was also shown by a classical treatment;^{7,15} (3) by eq. (5) and its complex conjugate, we find that the equation of motion for $B_j^* B_j$ contains no T_2 dephasing term, i.e., no vibrational energy is exchanged among the intramolecular modes due to the T_2 dephasing processes; (4) the amplitude function is related to the average excitation in the Poisson distribution¹⁶

$$P_{n_j}(t) = \exp(-\bar{n}_j) \bar{n}_j^{n_j} / n_j! , \quad (7)$$

where P_{n_j} is the energy population of a harmonic oscillator (for $\lambda=0$) and n_j and \bar{n}_j are the vibrational quantum number and the average excitation of the j -th mode, respectively; and

$$\bar{n}_j(t) = |B_j(t)|^2 = \sum_{n_j} n_j P_{n_j}(t) . \quad (8)$$

Therefore the average excitation $\bar{n}_j = |B_j|^2$ is a relevant quantity in describing the energy distribution among the intramolecular modes.

3. Dynamical Excitation and Optimal Detuning

(A) SF₆ gas-phase system

To show the role of the external and internal detunings in IVR, we now first consider the gas-phase SF₆ system in the collisionless low-pressure regime (i.e. $\Gamma_j=0$) and focus on the case of the fourth-order anharmonic coupling $\epsilon Q_3 Q_2^2 Q_6$, where ϵ is the anharmonicity and Q_3, Q_2 and Q_6 are the normal coordinates of the ν_3 (active), ν_2 and ν_6 modes. The corresponding internal detuning then is $\Delta_I = 2\pi(\nu_3 - 2\nu_2 - \nu_6)$ and the external detuning is $\Delta_E = 2\pi\nu_3 - \omega$,

where ω is the frequency of the tunable CO_2 TEA laser which is near resonant to the ν_3 mode. The quantum coupled equations of motion [eq. (7)] for this case become

$$i\hbar \dot{B}_1 = \lambda' C_1^4 B_1^* B_1^2 + \lambda' C_1 C_2^2 C_3 B_2^2 B_3^* \exp(i\Delta'_1 t) + (V_0/2) C_1 \exp(i\Delta'_E t), \quad (9.a)$$

$$i\hbar \dot{B}_2 = \lambda' C_2^4 B_2^* B_2^2 + 2\lambda' C_1 C_2^2 C_3 B_1 B_3^* \exp(i\Delta'_1 t), \quad (9.b)$$

$$i\hbar \dot{B}_3 = \lambda' C_3^4 B_3^* B_3^2 + \lambda' C_1 C_2^2 C_3 B_1^* B_2^2 \exp(i\Delta'_E t), \quad (9.c)$$

where B_j ($j=1,2,3$) are the related amplitude functions of the normal modes ν_3 , ν_2 and ν_6 , respectively; $\lambda'=12\lambda$, $\Delta'_1=2\pi(\nu'_3 + \nu'_6 - 2\nu'_2)$ and $\Delta'_E=2\pi\nu'_3 - \omega$, with ν'_j ($j=2,3,6$) are the anharmonic-corrected frequencies, i.e., $\nu'_j = \nu_j - \lambda' C_j / \hbar$; C_j are the quantization constants $C_j = (\hbar / 2m_j \omega_j)^{1/2}$. The above system, for the case of exact resonance $\Delta'_E=0$, was discussed and the related IVR was estimated by Gan et al.¹² Here we shall show the dynamical features of IVR and find the optimal external and internal detunings by solving the coupled equations of motion numerically. We investigate the time evolution of the average excitations, $\bar{n}_1 = |B_1|^2$, $\bar{n}_2 = |B_2|^2$ and $\bar{n}_3 = |B_3|^2$, governed by eqs. (9) with the initial Boltzmann populations at room temperature ($T=300\text{K}$), $B_1(0)=0.10$, $B_2(0)=0.214$ and $B_3(0)=0.417$, and the initial phases $\tan^{-1}[I_j(0)/R_j(0)]=0$, where I_j and R_j are the imaginary and real parts of B_j ($j=1,2,3$).

Fig. 1 shows the peak value of the average excitation (\bar{n}_1^*) of the active (ν_3) mode as a function of the external detuning with laser intensity $I=10^8 \text{ W/cm}^2$. It is seen that the optimal external detuning $\Delta_E^* = 17 \text{ cm}^{-1}$ and the value of \bar{n}_1^* decreases rapidly for Δ_E' larger than Δ_E^* . This

"red-shift" behavior of the optimal detuning is a general feature of any laser-excited anharmonic oscillator where the energy spacings are closer for higher levels. [For more detailed analysis we refer the reader to Ref. 17.] Figs. 2(A) and (B) show the exact resonance and the far-off resonance, respectively. It is seen that the peak value of the ν_2 mode (\bar{n}_2^*) decreases from $\bar{n}_2^* \approx 0.4$ (for $\Delta'_I = 0$) to $\bar{n}_2^* \approx 0.1$ (for $\Delta'_I = 63 \text{ cm}^{-1}$). By our numerical results, we estimate that the rising times of the average excitations t_1^* [defined by the first peak values of the average excitation $\bar{n}_1^*(t^*)$] range from $t_1^* = 27$ picoseconds (ps) to $t_1^* = 70$ ps depending on the internal detuning Δ'_I . However, the rising times of the ν_2 mode, i.e., the IVR times are in the order of $t_2^* \approx 10$ ps (or 3.3 cm^{-1}). A typical excitation diagram is shown in Fig. 2(C) for the case of $(\Delta'_E/2\pi, \Delta'_I/2\pi) = (2.71, 1.00)$. Finally, we consider the optimal internal detuning where the intramolecular anharmonic coupling is the most significant (or the maximum excitation of the ν_2 mode \bar{n}_2^*). We see, from Fig. 3, that the optimal internal detuning is $\Delta'_I/2\pi \approx 1.6 \text{ cm}^{-1}$ when the external detuning is fixed at its optimal value $\Delta'_E/2\pi \approx 2.7 \text{ cm}^{-1}$.

(B) H/W heterogeneous system

The IR spectrum of H/W is given by (in units of cm^{-1})¹⁸: $\nu_1 = 1048.6$, $\nu_2 = 645.3$ and $\nu_3 = 1290.6$. The fourth-order anharmonic coupling is given by $\epsilon' Q_1^2 Q_2 Q_3$, i.e., $\Delta'_I = 2\pi(2\nu_1 - \nu_2 - \nu_3)$, where the ν_1 mode is vibrationally excited by a CO_2 laser. Again, we investigate the average excitations of these three modes for the case of $\Gamma_j = 0$. Since the surface-phonon-induced level width (Γ_j) simply causes the decay of the energy excitations, it does not affect the rates of IVR significantly.⁵

Two typical dynamical features of the average excitations \bar{n}_1 , \bar{n}_2 and \bar{n}_3 (for ν_1 , ν_2 and ν_3 modes, respectively) with laser intensity $I=10^8$ W/cm² and the anharmonicity $\epsilon'=5$ cm⁻¹ are shown in Figs. 4(A) and (B). It is seen that these dynamical patterns of the excitations governed by $\epsilon'Q_1^2Q_2Q_3$ are significantly different from that of the SF₆ system governed by $\epsilon Q_3Q_2^2Q_6$. We note that the former system involves the operator function $a_1^2a_2^+a_3^+$ corresponding to a two-photon processes followed by an anharmonic two-quanta coupling, while the latter system involves $a_3(a_2^+)^2a_6^+$ corresponding to a single-photon processes followed by an anharmonic three-quanta coupling. We also note that the IVR rate of the H/W system is faster than that of the SF₆ system; however, the average excitation of the active mode in H/W system is lower since the two-photon transition (governed by $\mu_0''EQ^2$) is usually less probable than the single-photon transition, i.e., the dipole transition governed by $\mu_0'EQ$, where μ_0' and μ_0'' are, respectively, the first and second derivatives of the active dipole moment with respect to the normal coordinate Q and E is the local electric field acting on the species.

In conclusion, we note that for a heterogeneous system the local electric field acting on the adspecies may be greatly enhanced by the surface effects such as roughness.¹⁹ Furthermore, the surface bond breaking can also be possible via van der Waals interaction or the surface-migration-induced rate processes.^{20,21}

One of us (J.L.) would like to thank Prof. Z. Gan of Peking University for useful communications and sending their Chinese articles on SF₆.

*This work was supported in part by the Office of Naval Research, the Air Force Office of Scientific Research (AFSC), United States Air Force, under Grant AFOSR-82-0046 and the National Science Foundation under Grant No. CHE-8022874.

†Camille and Henry Dreyfus Teacher-Scholar (1975-82).

1. For recent reviews, see N. Bloembergen and E. Yablonovitch, *Physics Today* 31, 23 (1978), and a Special Issue on Laser Chemistry, *Physics Today* 33, 25 (1980).
2. J. Lin, *Phys. Lett.* 70A, 195 (1979).
3. J. Lin and T. F. George, *Phys. Lett.* 80A, 296 (1980).
4. J. Lin and T. F. George, *Surf. Sci.* 100, 381 (1980).
5. J. Lin and T. F. George, *J. Phys. Chem.* 84, 2957 (1980).
6. J. Lin and T. F. George, *Phys. Rev. B*, (1981) in press.
7. J. Lin and T. R. George, *Surf. Sci.* 107, 417 (1981).
8. J. Lin and T. F. George, *Chem. Phys. Lett.* 66, 5 (1979).
9. J. Lin and T. F. George, *J. Chem. Phys.* 72, 2554 (1980).
10. J. Lin, Ph.D. Thesis, University of Rochester (1980).
11. J. Lin and T. F. George, *Surf. Sci.*, (1981) in press.
12. Z. Gan, G. Yang, X. Huang and K. Feng, *ACTA PHYSICA SINICA* (Peking) 29, 743 (1980).
13. J. Lin, A. C. Beri, M. Hutchinson, W. C. Murphy and T. F. George, *Phys. Lett.* 79A, 233 (1980).
14. W. H. Louisell, Quantum Statistical Properties of Radiation (Wiley, New York, 1973).
15. J. Ford, *J. Math. Phys.* 2, 387 (1961).
16. E. Merzbacher, Quantum Mechanics, 2nd ed. (Wiley, New York, 1970), p. 366.

17. T. F. George, J. Lin, K. S. Lam and C. Chang, Opt. Eng. 19, 100 (1980).
18. R. F. Willis, W. Ho and E. W. Plummer, Surf. Sci. 80, 593 (1979).
19. R. P. Van Duyne, in: Chemical and Biochemical Applications of Lasers, Vol. IV, ed. by C. B. Moore (Academic Press, New York, (1979), Chap. 4.
20. D. Lucas and G. E. Ewing, private communication.
21. J. Lin and T. F. George, to be published.

Figure Captions

- Fig. 1. The peak value of the average excitation of the active mode (\bar{n}_1^*) vs. the external detuning (Δ'_E). [SF_6 system].
- Fig. 2. The time-dependent average excitations of the active mode (\bar{n}_1), the ν_2 mode (\bar{n}_2) and the ν_6 mode (\bar{n}_3) for the detunings $(\Delta'_E/2\pi, \Delta'_I/2\pi) =$ (A) (0,0) exact resonance, (B) (0,20) far-off internal resonance and (C) (2.71,1.00). $\Delta'_E/2\pi = 2.71$ is the optimal external detuning.
- Fig. 3. The peak value of the average excitation of the ν_2 mode (\bar{n}_2^*) vs. the internal detuning (Δ'_I) with the optimal external detuning $\Delta'_E = \Delta_E^*$.
- Fig. 4. The dynamical features of the average excitations of the active mode (\bar{n}_1), the ν_2 mode (\bar{n}_2) and the ν_3 mode (\bar{n}_3) for a heterogeneous system H/W with detunings $(\Delta'_E/2\pi, \Delta'_I/2\pi) =$ (A) (0,0) exact resonance and (B) (10,80).





Fig. 1






Fig. 3

AVERAGE EXCITATIONS





Fig. 4

AVERAGE EXCITATIONS



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, N.C. 27709	1
ONR Western Regional Office Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Eastern Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Commander, Naval Air Systems Command Attn: Code 3100 (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. David L. Nelson Chemistry Program Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720	1	Dr. C. P. Flynn Department of Physics University of Illinois Urbana, Illinois 61801	1
Dr. L. N. Jarvis Surface Chemistry Division 4555 Overlook Avenue, S.W. Washington, D.C. 20375	1	Dr. W. Kohn Department of Physics University of California (San Diego) LaJolla, California 92037	1
Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181	1	Dr. R. L. Park Director, Center of Materials Research University of Maryland College Park, Maryland 20742	1
Dr. John T. Yates Department of Chemistry University of Pittsburgh Pittsburgh, Pennsylvania 15260	1	Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455	1
Dr. Theodore E. Madey Surface Chemistry Section Department of Commerce National Bureau of Standards Washington, D.C. 20234	1	Dr. Chia-wei Woo Department of Physics Northwestern University Evanston, Illinois 60201	1
Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712	1	Dr. D. C. Mattis Polytechnic Institute of New York 333 Jay Street Brooklyn, New York 11201	1
Dr. Keith H. Johnson Department of Metallurgy and Materials Science Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. Robert M. Hexter Department of Chemistry University of Minnesota Minneapolis, Minnesota 55455	1
Dr. J. E. Demuth IBM Corporation Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598	1	Dr. R. P. Van Duyne Chemistry Department Northwestern University Evanston, Illinois 60201	1

TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. S. Sibener Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637	1	Dr. Martin Fleischmann Department of Chemistry Southampton University Southampton SO9 5NH Hampshire, England	1
Dr. M. G. Lagally Department of Metallurgical and Mining Engineering University of Wisconsin Madison, Wisconsin 53706	1	Dr. J. Osteryoung Chemistry Department State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Robert Gomer Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637	1	Dr. G. Rubloff I.B.M. Thomas J. Watson Research Center P. O. Box 218 Yorktown Heights, New York 10598	1
Dr. R. G. Wallis Department of Physics University of California, Irvine Irvine, California 92664	1	Dr. J. A. Gardner Department of Physics Oregon State University Corvallis, Oregon 97331	1
Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052	1	Dr. G. D. Stein Mechanical Engineering Department Northwestern University Evanston, Illinois 60201	1
Dr. P. Hansma Chemistry Department University of California, Santa Barbara Santa Barbara, California 93106	1	Dr. K. G. Spears Chemistry Department Northwestern University Evanston, Illinois 60201	1
Dr. P. Hendra Chemistry Department Southampton University England SO9JNH	1	Dr. R. W. Plummer University of Pennsylvania Department of Physics Philadelphia, Pennsylvania 19104	1
Professor P. Skell Chemistry Department Pennsylvania State University University Park, Pennsylvania 16802	1	Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	2
Dr. J. C. Hemminger Chemistry Department University of California, Irvine Irvine, California 92717	1	Professor D. Hercules University of Pittsburgh Chemistry Department Pittsburgh, Pennsylvania 15260	1

TECHNICAL REPORT DISTRIBUTION LIST, 056

No.
Copies

Professor N. Winograd
The Pennsylvania State University
Department of Chemistry
University Park, Pennsylvania 16802 1

Professor J. F. George
The University of Rochester
Chemistry Department
Rochester, New York 14627 1

Professor Dudley R. Herschbach
Harvard College
Office for Research Contracts
1350 Massachusetts Avenue
Cambridge, Massachusetts 02138 1

Professor Horia Metiu
University of California,
Santa Barbara
Chemistry Department
Santa Barbara, California 93106 1

Professor A. Steckl
Rensselaer Polytechnic Institute
Department of Electrical and
Systems Engineering
Integrated Circuits Laboratories
Troy, New York 12181 1

Professor R. D. Archer
University of Massachusetts
Chemistry Department
Amherst, Massachusetts 01003 1

Dr. A. C. Pastor
Hughes Research Laboratories
3011 Malibu Canyon Road
Malibu, California 90265 1

DATE
ILME